arXiv:1410.5002v1 [math.ST] 18 Oct 2014

UNBIASED ESTIMATORS FOR ENTROPY AND CLASS NUMBER

STEPHEN MONTGOMERY-SMITH AND THOMAS SCHÜRMANN

Abstract. We introduce unbiased estimators for the Shannon entropy and the class number, in the situation that we are able to take sequences of independent samples of arbitrary length.

Introduction. This paper supposes that we may pick a sequence of arbitrary length of independent samples $w_1, w_2, \dots$ from an infinite population. Each sample belongs to one of M classes $C_1, C_2, \dots, C_M$, and the probability that a sample belongs to class C_i is p_i. So these probabilities satisfy the constraints $0 \le p_i \le 1$ and $\sum_{i=1}^M p_i = 1$.

The goal of this paper is to present methods to estimate the Shannon entropy $H = -\sum_{i=1}^M p_i \log(p_i)$ (see [8]). An obvious method is to take a sample of size n, and compute the estimators $\hat{p}_i = k_i/n$, where k_i are the number of samples from the class C_i. However this is known to systematically underestimate the entropy, and it can be significantly biased [2][4][5][6]. Recently there have been more advanced estimators for the entropy which have smaller bias [2][3][4][5][6][7].

In this paper we introduce new entropy estimators that have bias identically zero. We also introduce an unbiased estimator for the class number M, a problem of interest to ecologists (see for example the review article [1]). The disadvantage of all our methods is that there is no *a priori* estimate of the sample size. For this reason, we postpone rigorous analysis of variance and other measures of confidence until it becomes clear that these estimators are of more than theoretical value.

We will use the following power series. Define the harmonic number by $h_n = \sum_{k=1}^n 1/k$, $h_0 = 0$. Then for $|x| < 1$

$$\frac{1}{(1-x)^2} = \sum_{k=1}^\infty k x^{k-1},$$

$$\log(1-x) = -\sum_{k=1}^\infty \frac{x^k}{k},$$

$$\frac{\log(1-x)}{(1-x)} = -\sum_{k=1}^\infty h_k x^k.$$

First Estimator for Entropy. For each $1 \le i \le M$, let N_i denote the smallest $k \ge 1$ for which $w_k \in C_i$. Then

$$\hat{H}_1 = \sum_{i=1}^M \frac{I_{N_i \ge 2}}{N_i - 1}$$

is an unbiased estimator for the entropy. The proof is straightforward. The marginal distribution of N_i satisfies the geometric distribution $\Pr(N_i = k) = p_i(1-p_i)^{k-1}$. Thus

$$E\left(\frac{I_{N_i \geq 2}}{N_i - 1}\right) = \sum_{k=2}^{\infty} \frac{p_i(1-p_i)^{k-1}}{k-1} = -p_i \log(p_i).$$

Depending upon the applications, a possible disadvantage of this estimator is that complete knowledge of all possible classes needs to be known in advance.

Second Estimator for Entropy. Let N denote the smallest $k \geq 1$ such that w_1 and w_{k+1} belong to the same class. Then $\hat{H}_2 = h_{N-1}$ is an unbiased estimator for the entropy.

This follows, since conditional upon $w_1 \in C_i$, the distribution of N satisfies the geometric distribution $\Pr(N = k | w_1 \in C_i) = p_i(1-p_i)^{k-1}$. Thus

$$E(h_{N-1} | w_1 \in C_i) = \sum_{k=1}^{\infty} h_{k-1} p_i (1-p_i)^{k-1} = -\log(p_i),$$

and hence

$$E(h_{N-1}) = \sum_{i=1}^{M} E(h_{N-1} | w_1 \in C_i) \Pr(w_1 \in C_i) = -\sum_{i=1}^{M} p_i \log(p_i).$$

While we don't wish to focus on analysis of the variance, it is certainly clear that this single estimator by itself will have unusable confidence limits. While the variance can be reduced by taking the mean of n of these estimators, we propose the following version. For each $1 \leq j \leq n$, let $N^{(j)}$ be the smallest $k \geq 1$ such that w_j and w_{k+j} belong to the same class. Then the unbiased estimator is

$$\hat{H}_3 = \frac{1}{n} \sum_{j=1}^{n} h_{N^{(j)}-1}.$$

Estimator for Class Number. This is very similar to the second estimator for entropy. Define N and $N^{(j)}$ as in the previous section. Then $\hat{M}_1 = N$ is an unbiased estimator for the class number M. The proof is almost identical to that provided in the previous section.

We may also produce an unbiased estimator

$$\hat{M}_2 = \frac{1}{n} \sum_{j=1}^{n} N^{(j)}.$$

This last quantity can also be considered as a corrector to the naive estimator $\hat{M}_3$, which is defined as the number of classes observed in the first n samples, or alternatively, as the cardinality of the set A, where A is the set of $1 \leq i \leq M$ such that there exists $1 \leq k \leq n$ for which $w_k \in C_i$.

After picking n samples, then continue picking samples until every class observed in the first n samples is observed at least once more. For each $i \in A$, let F_i denote the smallest $k \geq 1$ such that $w_k \in C_i$, and let L_i denote the smallest $k \geq 1$ such that $w_{k+n} \in C_i$. Then

$$\hat{M}_2 = \hat{M}_3 + \frac{1}{n} \sum_{i \in A} (L_i - F_i).$$

If one doesn't wish to record the order in which the samples are obtained, one can simply compute the expected value of this quantity over all possible rearrangements of obtaining this data, and derive the unbiased estimator

$$\hat{M}_4 = \hat{M}_3 + \frac{1}{n}\sum_{i\in A}\left(\frac{m+1}{s_i+1} - \frac{n+1}{r_i+1}\right),$$

where r_i is the number of times the ith class appears in the first sample of size n, m is the size of the subsequent sample, and s_i is the number of times the ith class appears in the subsequent sample.

References

[1] A. Chao, Species richness estimation. In N. Balakrishnan, C. B. Read, and B. Vidakovic, eds., *Encyclopedia of Statistical Sciences* pp 7909-7916 (New York, Wiley, 2005).

[2] P. Grassberger, Phys. Lett. **A 128**, 369 (1988).

[3] P. Grassberger, www.arxiv.org, physics/0307138 (2003).

[4] B. Harris, The statistical estimation of entropy in the non-parametric case *Topics in Information Theory* ed I Csiszar (Amsterdam: North-Holland) pp 323-55 (1975).

[5] H. Herzel, Sys. Anal. Mod. Sim. **5**, 435 (1988).

[6] G. Miller, Note on the bias of information estimates. In H. Quastler, ed., *Information theory in psychology II-B*, pp 95-100 (Free Press, Glencoe, IL 1955).

[7] T. Schürmann, J. Phys. A: Math. Gen. **37**, L295-L301 (2004).

[8] C. E. Shannon and W. Weaver, *The Mathematical Theory of Communication*, (University of Illinois Press, Urbana, IL 1949).

Math Dept, University of Missouri, Columbia, MO 65211, U.S.A.
E-mail address: stephen@missouri.edu

Jülich Supercomputing Centre, Jülich Research Centre, 52425 Jülich, Germany.
E-mail address: thomas.schuermann@live.de